




# Quick guide for fertilizing **transplanted rice** in **Ilocos Sur** (straw removed)



**Fertilizer application scheme (bags/ha)** 1 cavan = 50 kg

Step no.	Growth stage	Days after transplanting (DAT)			Fertilizer source	Yield target (based on total fresh weight from the thresher)				
		Early maturing	Medium early maturing	Medium maturing		80-100 cavans/ha	100-125 cavans/ha	125-150 cavans/ha	150-170 cavans/ha	
									Option #1	Option #2
Step 1	Early growth	0 - 14	0 - 14	0 - 14	14-14-14 (with S)	2 bags	3 bags	4 bags	5 bags	5 bags
Step 2	Active tillering	23 - 27	24 - 28	28 - 32	Urea (46-0-0)	1 bag	1.5 bags	2 bags	2 bags	2 bags
Step 3	Panicle initiation	33 - 37	35 - 39	43 - 47	Urea (46-0-0)	1 bag	1.5 bags	2 bags	2.5 bags	1.5 bags
					MOP (0-0-60)				0.5 bag	
					17-0-17 (with S)					2 bags

This 'quick guide' is applicable for rice fields with:

- Sufficient supply of water to avoid drought
- An average level of soil fertility
- Use of seedlings less than 23 days old
- No input of nutrient through deposition of sediment from flooding
- No use of organic materials other than rice stubble

For fields with other conditions, use 'Nutrient Manager for Rice' to develop a field-specific guideline. Consult the local DA office, PhilRice, ATI, Office of the Provincial Agriculturist (OPAg), and MAO in your municipality about use of the 'Nutrient Manager for Rice'.

This is a joint undertaking of DA, PhilRice, OPAg, and IRRI