




Quick guide for fertilizing **transplanted rice** in Negros Oriental (**straw removed**)



Fertilizer application scheme (bags/ha)

1 cavan = 50 kg

Step no.	Growth stage	Days after transplanting (DAT)			Fertilizer source	Yield target (based on total fresh weight from the thresher)			
		Early maturing	Medium early maturing	Medium maturing		60-80 cavans/ha	80-100 cavans/ha	100-125 cavans/ha	125-150 cavans/ha
Step 1	Early growth	0 – 14	0 – 14	0 – 14	16-20-0 (with S)	1 bag			
					14-14-14 (with S)		2 bags	3 bags	4 bags
Step 2	Active tillering	23 – 27	24 – 28	28 – 32	Urea (46-0-0)		1 bag	1.5 bags	2 bags
Step 3	Panicle initiation	33 – 37	35 – 39	43 – 47	Urea (46-0-0)	1 bag	1 bag	1.5 bags	2 bags

This 'quick guide' is applicable for rice fields with:

- Sufficient supply of water to avoid drought
- Use of seedlings less than 23 days old
- No input of nutrient through deposition of sediment from flooding
- No use of organic materials other than rice stubble

For fields with other conditions, use 'Nutrient Manager for Rice' to develop a field-specific guideline. Consult the local DA office, PhilRice, ATI, Office of the Provincial Agriculturist (OPAg), and Agricultural Technicians in your municipality about use of the 'Nutrient Manager for Rice'.

This is a joint undertaking of DA, PhilRice, OPAG, and IRRI

Version 1.0 July 2009